arXiv:1211.6413v1 [math.PR] 27 Nov 2012

Preprint submitted to arXiv (28 November 2012)

A LIMIT THEOREM FOR THE SUM OF SQUARED DIFFERENCES OF AN INTEGRATED ITO PROCESS WITH APPLICATION TO INVERSE SCATTERING

JOHN F. A. FLETCHER,* *University of Cambridge*

Abstract

We investigate a functional obtained by summing the squared differences of the integral of an Ito process over disjoint intervals. The limit of this sum is shown to converge in probability to two thirds the quadratic variation of the underlying process. An application to inverse scattering from a random fractal surface is presented.

Keywords: Quadratic variation; local average; inverse scattering

2010 Mathematics Subject Classification: Primary 60F17
Secondary 78A46: 78A46

Introduction

Imagine that the real-valued function $(X_t)_{t\in\mathbb{R}}$ represents the density of a physical quantity. For example, X could be the density of a fluid along a one-dimensional subspace parametrised by s, or the energy density of radiation in a scattering problem. With regard to measurements, such densities are, of course, idealisations. In reality, one would attempt to *estimate* the value of X, for fixed s, by taking some sort of local average. Define a discrete set of times $t_k = \Delta k$ where $k = 0, 1, 2, ...$, and $\Delta > 0$. Denote the average of X over the k^{th} interval $[t_k, t_{k+1})$ by $\hat{X}_k$ where

$$\hat{X}_k := \frac{1}{\Delta} \int_{t_k}^{t_{k+1}} X_s \, \mathrm{d}s. \tag{1}$$

---

* Postal address: DAMTP, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA. Email address: j.f.a.fletcher@damtp.cam.ac.uk

1. Local average and quadratic variation

Quadratic variation is a useful notion in many circumstances, for example, it can provide a way to estimate the volatility of the quantity under study. If local averages of the type (1) are available, the following theorem may be applied:

Theorem 1.1. *For a standard Ito process* $(X_t)_{t\geq 0}$ *where*

$$X_t = X_0 + \int_0^t a_s \,\mathrm{d}s + \int_0^t b_s \,\mathrm{d}B_s \tag{2}$$

with average $\hat{X}_k$ *defined by* (1) *we have*

$$\sum_{k=1}^{n-1} (\hat{X}_k - \hat{X}_{k-1})^2 \overset{\mathbb{P}}{\to} \frac{2}{3}\langle X\rangle_t \quad \text{as} \quad \Delta \to 0 \tag{3}$$

where $\{t_k; k = 0, 1, ..., n = t/\Delta\}$ *specifies a partition of* $[0, t]$, *and* $\overset{\mathbb{P}}{\to}$ *denotes convergence in probability.*

Proof. Integrating X by parts:

$$\int\limits_{t_k}^{t_{k+1}} X_s \,\mathrm{d}s = sX_s\big|_{t_k}^{t_{k+1}} - \int\limits_{t_k}^{t_{k+1}} s \,\mathrm{d}X_s. \tag{4}$$

It follows that

$$\begin{aligned} \hat{X}_k - \hat{X}_{k-1} &= \frac{1}{\Delta} \int\limits_{t_{k-1}}^{t_k} (s - t_{k-1}) \,\mathrm{d}X_s + \frac{1}{\Delta} \int\limits_{t_k}^{t_{k+1}} (t_{k+1} - s) \,\mathrm{d}X_s \\ &= \int\limits_{t_{k-1}}^{t_{k+1}} \rho_s(k, \Delta) \,\mathrm{d}X_s \end{aligned}$$

where

$$\rho_s(k, \Delta) = \begin{cases} (s - t_{k-1})/\Delta & \text{if } t_{k-1} \leq s < t_k, \\ (t_{k+1} - s)/\Delta & \text{if } t_k \leq s < t_{k+1}. \end{cases}$$

Assume for definiteness that n is even (the case of odd n follows in an obvious way).

Splitting the sum in (3) into two:

$$\begin{aligned}
&\sum_{k=1}^{n-1}(\hat{X}_k - \hat{X}_{k-1})^2 \\
&= \sum_{k=1}^{n-1}\left(\int\limits_{t_{k-1}}^{t_{k+1}} \rho_s(k,\Delta)\,\mathrm{d}X_s\right)^2 \\
&= \sum_{k=1}^{n/2}\left(\int\limits_{t_{2k-2}}^{t_{2k}} \rho_s(2k-1,\Delta)\,\mathrm{d}X_s\right)^2 + \sum_{k=1}^{n/2-1}\left(\int\limits_{t_{2k-1}}^{t_{2k+1}} \rho_{s-\Delta}(2k-1,\Delta)\,\mathrm{d}X_s\right)^2 . \qquad (5)
\end{aligned}$$

Define the continuous function $\eta_t(\Delta)$ as

$$\eta_t(\Delta) = \sum_{k=1}^{\infty} \mathbf{1}_{\{t_{2k-2}\le t<t_{2k}\}}\rho_t(2k-1,\Delta) \tag{6}$$

and the process $\widetilde{X}_t := X_0 + \int_0^t \eta_s(\Delta)\,\mathrm{d}X_s$. Also, we use the following notation

$$Q_\Delta^{(N)}(X_t) := \sum_{k=1}^{n} |X_{k\Delta} - X_{(k-1)\Delta}|^N. \tag{7}$$

Set $u_k := t_{2k}$. The first sum in (5) becomes

$$\sum_{k=1}^{n/2}\left(\int\limits_{t_{2k-2}}^{t_{2k}} \rho_s(2k-1,\Delta)\,\mathrm{d}X_s\right)^2 = \sum_{k=1}^{n/2}\left(\widetilde{X}_{u_k} - \widetilde{X}_{u_{k-1}}\right)^2. \tag{8}$$

The the limit $\Delta \to 0$ of (8) looks a lot like a quadratic variation, except that $\widetilde{X}$ is dependent on the grid spacing Δ.

The process X can be expressed as $X_t \equiv A_t + Z_t$ where A has finite variation and Z is a local martingale. Clearly $|\widetilde{Z}_{u_k} - \widetilde{Z}_{u_{k-1}}| \le |Z_{u_k} - Z_{u_{k-1}}|$ and $|\widetilde{A}_{u_k} - \widetilde{A}_{u_{k-1}}| \le |A_{u_k} - A_{u_{k-1}}|$ for all k. As one might expect, given the well known properties of the quadratic variation, the process $\widetilde{A}$ does not contribute to (8) in the limit $\Delta \to 0$. This can be seen in the following way:

$$\begin{aligned}
\Big|\sum_k(\widetilde{Z}_{u_k} - \widetilde{Z}_{u_{k-1}})(\widetilde{A}_{u_k} - \widetilde{A}_{u_{k-1}})\Big| &\le \left(\sup_k |\widetilde{Z}_{u_k} - \widetilde{Z}_{u_{k-1}}|\right)\cdot Q_{2\Delta}^{(1)}(\widetilde{A}_t) \\
&\le \left(\sup_k |Z_{u_k} - Z_{u_{k-1}}|\right)\cdot Q_{2\Delta}^{(1)}(A_t)
\end{aligned}$$

Compare proposition 1.18, p.128, [3].

and this converges to zero as $\Delta \to 0$ because of the continuity of Z. Similarly

$$\lim_{\Delta \to 0} \sum_k (\widetilde{A}_{u_k} - \widetilde{A}_{u_{k-1}})^2 = 0. \tag{9}$$

We now require two well known results from the literature, which are stated as the following Lemmas:

Lemma 1.1. *Let* $(M_s)_{0 \leq s \leq t}$ *be a uniformly continuous martingale such that* $|M_s| \leq K < \infty$, $\mathbb{P}$-*a.s. then*

$$\lim_{\Delta \to 0} \mathbb{E}[Q_\Delta^{(4)}(M_t)] = 0.$$

Proof. See for example Lemma 5.10, p.33, [1].

Lemma 1.2. *Let* $\langle M \rangle_s \leq K$ *for all* $s \in [0, t]$, $\mathbb{P}$-*a.s. then*

$$\mathbb{E}\left[\left(Q_\Delta^{(2)}(M_t) - \langle M \rangle_t\right)^2\right] \leq 2\mathbb{E}\big[Q_\Delta^{(4)}(M_t)\big] + 2\mathbb{E}\big[\langle M \rangle_t \cdot m_t\left(\Delta, \langle M \rangle\right)\big]$$

where $m_u(\Delta, X) := \sup\{|X_t - X_s|; 0 \leq s < t \leq u \text{ and } |t - s| \leq \Delta\}$ *is the modulus of continuity.*

Proof. See proof of theorem 5.8, p.34, [1].

Assume that Z is a bounded, uniformly continuous martingale. Lemma 1.1 implies

$$\mathbb{E}[Q_{2\Delta}^{(4)}(\widetilde{Z}_t)] \leq \mathbb{E}[Q_{2\Delta}^{(4)}(Z_t)] \to 0 \quad \text{as} \quad \Delta \to 0. \tag{10}$$

Furthermore

$$m_t(2\Delta, \langle \widetilde{Z} \rangle) \leq m_t(2\Delta, \langle Z \rangle) \to 0 \quad \text{as} \quad \Delta \to 0 \tag{11}$$

by the sample path uniform continuity of $\langle Z \rangle$. Assume that $\langle Z \rangle$ is bounded, then so is $\langle \widetilde{Z} \rangle$, and, since $m_t(2\Delta, \langle \widetilde{Z} \rangle)$ is bounded

$$\lim_{\Delta \to 0} \mathbb{E}\big[\langle \widetilde{Z} \rangle_t \cdot m_t(2\Delta, \langle \widetilde{Z} \rangle)\big] = 0 \tag{12}$$

by the dominated convergence theorem and (11). Putting these facts together and using Lemma 1.2 we have

$$\lim_{\Delta \to 0} \mathbb{E}\left[\left(Q_{2\Delta}^{(2)}(\widetilde{Z}_t) - \langle \widetilde{Z} \rangle_t\right)^2\right] = 0. \tag{13}$$

Convergence in L^2 implies convergence in probability.

Next, we would like to show that $\langle \widetilde{Z} \rangle$ converges, in some sense, to a well defined limit, independent of Δ.

$$\begin{aligned}\langle \widetilde{Z} \rangle_t =& \sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \eta_s^2(\Delta) b_s^2 \,\mathrm{d}s \\ =& \sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \eta_s^2(\Delta) \left(b_s^2 - b_{u_k^*}^2 \right) \mathrm{d}s + \sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \eta_s^2(\Delta) b_{u_k^*}^2 \,\mathrm{d}s \end{aligned} \tag{14}$$

where $u_k^* \in [u_{k-1}, u_k)$ is such that $b_{u_k^*} = \sup\{b_s; u_{k-1} \leq s \leq u_k\}$. The first integral on the right hand side of (14) has the bound

$$\begin{aligned}\sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \eta_s^2(\Delta) \left(b_s^2 - b_{u_k^*}^2 \right) \mathrm{d}s \leq& \sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \left(b_s^2 - b_{u_k^*}^2 \right) \mathrm{d}s \\ =& \int_0^t b_s^2 \,\mathrm{d}s - \sum_{k=1}^{n/2} 2\Delta b_{u_k^*}^2 \xrightarrow{\text{a.s.}} 0 \quad \text{as} \quad \Delta \to 0.\end{aligned}$$

For the second integral in (14) we have

$$\begin{aligned}&\sum_{k=1}^{n/2} \int\limits_{u_{k-1}}^{u_k} \eta_s^2(\Delta) b_{u_k^*}^2 \,\mathrm{d}s \\ &= \frac{1}{3} \sum_{k=1}^{n/2} 2\Delta b_{u_k^*}^2 \xrightarrow{\text{a.s.}} \frac{1}{3} \int_0^t b_s^2 \,\mathrm{d}s \quad \text{as} \quad \Delta \to 0\end{aligned} \tag{15}$$

and so

$$Q_{2\Delta}^{(2)}(\widetilde{Z}_t) \xrightarrow{\mathbb{P}} \frac{1}{3} \langle Z \rangle_t \quad \text{as} \quad \Delta \to 0. \tag{16}$$

The technique of localisation can be used to generalise (16) to the case where Z and $\langle Z \rangle$ are not necessarily bounded, using the stopping time

$$T_j = \inf\{t \geq 0; |Z_t| \geq j \text{ or } \langle Z \rangle_t \geq j\}. \tag{17}$$

The limit of the second sum in (5) can be evaluated in precisely the same way as the first, and converges to the same limit. Finally (3) follows.

2. Application to inverse scattering

Norris [2] showed that, in the ray-theory limit, fluctuations in the intensity of light scattered from a corrugated surface with Brownian motion slope $(\sigma W_s)_{s \in \mathbb{R}}$ are described by the Ito process $(R_t)_{t\in\mathbb{R}}$ where

$$dR_t = \frac{2}{\sigma^2}(1 - R_t)dt + \frac{2}{\sigma}\sqrt{R_t}dB_t \tag{18}$$

and σ is a positive constant. Measurements of the average energy flux between points $\{t_k; u = t_0, t_1, ..., t_n = v\}$, such that $t_k - t_{k-1} = \Delta > 0$, correspond to $\hat{R}_k$. The volatility of the surface slope can be obtained approximately, for small Δ, from (3) since

$$\lim_{\Delta\to 0}\sum_{k=1}^{n-1}(\hat{R}_k - \hat{R}_{k-1})^2 = \frac{2}{3}\cdot\frac{4}{\sigma^2}\int_u^v R_t\,\mathrm{d}t = \frac{2}{3}\cdot\frac{4}{\sigma^2}\sum_{k=0}^{n-1}\hat{R}_k\Delta \tag{19}$$

convergence in probability.

Acknowledgements

The author would like to thank James Norris, Chris Rogers, Kalvis Jansons and Moritz Duembgen for useful discussions relating to this work.

References

[1] Karatzas, I. and Shreve, S. (1991). *Brownian motion and stochastic calculus* vol. 113. Springer Verlag.

[2] Norris, J. (1988). Intensity fluctuations of randomly scattered waves considered as local time. *Physics Letters A* **128,** 404–405.

[3] Revuz, D. and Yor, M. (1999). *Continuous martingales and Brownian motion* vol. 293. Springer Verlag.